

Ethiopian Meteorology Institute

Health-Meteorology Bulletin

The product of
Biometeorology and
insurance desk

Dekadal Bulletin

Volume: 1

Issued data: 01/02/2025

Valid until : 20/03/2025

February 2025

©*Ethiopian Meteorology
Institute*

Tell: 251(0)11 6615779

Fax: 251(0)11 6625292

P.O.Box 1090

Website:

<http://www.ethiomet.gov.et/>

Telegram:

<https://t.me/BiometService>



Table of Contents

I.	Foreword	3
II.	Assumptions	4
i.	Malaria	4
ii.	Human heat index:	4
iii.	Cattle heat index:	4
1.	Weather impact Assessment on Health for February 2025 First dekad	5
1.1	Malaria prone areas during February 2025 First dekad.....	5
1.2	Climate comfort Conditions	5
1.2.1	Comfortability for Human.....	5
1.2.2	Comfortability for Cattle.....	5
2.	Expected Weather Impact on Health for first dekad of February 2025 Second dekad	6
2.1	Expected Mosquito breeding areas.....	6
2.2	Temperature Humidity Index	6
2.2.1	Comfortability for Human	6
2.2.2	Comfortability for Cattle	6
3.	Summary	7
4.	Advisory	7

I. Foreword

This "Climate Information for the Health Sector" Bulletin has been designed to convey essential information regarding the monitoring of human comfort conditions based on the analysis of temperature and humidity data and also for the monitoring of Malaria outbreak areas based on the analysis of temperature and precipitation data. Since the monitoring of temperature and rainfall over a given area can be used to assess the likelihood of outbreak of Malaria with a lag of two months, this information can be an important for early warning tool if used judiciously.

The major objective of this bulletin is in line with the Ethiopia Meteorological Institute strategy of diversifying climate application products to the basic developmental sectors (such as the Health, the water, the agricultural sector etc....). This bulletin can be a very important source of information to Health professionals engaged in the monitoring of Public Health, to Tourism Agents and institutions who advise tourists regarding the comfort conditions of the places to be visited by the tourists and to the researcher who is interested in the field of Bio-Climatology.

We have the opinion that careful and continuous use of this bulletin can benefit to the improvement of early warning and preparedness in the Health sector.

Meanwhile, your comments and constructive suggestions are highly appreciated to make the objective of this bulletin a success,

This same bulletin can be accessed online at: http://www.ethiomet.gov.et/bulletins/health_bulletins

Director General
Ethiopia Meteorology Institute
P.O.Box 1090
Tel: 251(0)11 6615779
FAX 251(0)11 6625292
Web: www.ethiomet.gov.et
Addis Ababa, Ethiopia

II. Assumptions

i. Malaria: According to the International Research Institute for Climate and Society, (IRI), the predicted conditions of rainfall, temperature, and relative humidity are used in determining the degree of incidence for malaria.

- When rainfall is above 80 mm, the temperature is between 25°C and 32°C, and relative humidity is greater than 80%, the region is at high risk and is placed under high incidence.
- When the temperature is between 20°C and 25°C, relative humidity is between 70 and 80%, and rainfall is above 80 mm, then moderate incidence is advised.
- Low incidence for malaria is issued when the temperature is in the range of 18°C-20°C, relative humidity is 60 - 70% and rainfall is above 80 mm.
- No incidence is required when the temperature is less than 18°C, relative humidity less than 60%, and rainfall amount below 80 mm.

Based on these, climate variables have **a one to two months** postponed (delayed) effect on the spread of malaria.

ii. Human heat index: is a measure of how hot it feels when relative humidity is factored with the actual air temperature. The levels of caution for heat index are classified as follows:

- Cold stress when THI is <14, *Asthma, Pneumonia, Common Cold and flu*
- Comfortable when THI is 14-21, *pleasanter*
- Moderate when THI is 21-26, *No more effects*
- Heat stress when THI is >26, *heat stroke, heat cramps, hyperthermia, respiratory and cardiovascular diseases*

iii. Cattle heat index: The climatic condition for Cattle is a measure that accounts for the combined effects of environmental temperature and relative humidity on cattle. The level of heat stress for cattle classified as follows:

- Not Stressed when THI is <68, *free from heat stress*
- Stressed threshold when THI is 68 – 71, *impact less stress starting*
- Mild stress when THI is 72 – 79, *stress begins and calf rate affected*
- Moderate stress when THI is 80 – 89, *Milk production affected*
- Severe stress when THI is 90 – 99, *very significant losses in milk production*
- Extremely stress when THI is >100, *ultimate dead of cows*

1. Weather impact Assessment on Health for February 2025 First Dekad



1.1 Malaria prone areas during Feb1stDekad

During Feb1stDekad climate assessment for malaria breeding and transmission, in any parts of the country there was **no** suitable areas for malaria transmission over the country as illustrated in Figure 1




Figure 1: Malaria Prone areas

1.2 Climate comfort Conditions

1.2.1 Comfortability for Human

During the 1st dekad of February, the climate for human daily activities was very pleasant in most parts of the country. However, there was heat stress in the Gambella region and the working hours in the region were revised.as shown in figure 2.




Figure 2: Human comfort index

1.2.2 Comfortability for Cattle

Similar to the human's comfortability condition, there were no-significant heat stress climate conditions that impact the cattle's productivity, But mild heat stress were experienced in a few place of Southern Somali ,South west Ethiopia, Central Ethiopia and Gambela regions of the country as shown in figure 3.




Figure 3: Cattle Comfort index

2. Expected Weather Impact on Health for Second dekad of February 2025

2.1 Expected Mosquito breeding areas

In the coming Second Ten days of February 2025, the climate condition for malaria breeding and transmission **will not suitable** over most parts of the country except Tips area of North east Tigray as illustrated as red in figur4.




Figure 4: Expected malaria prone areas

2.2 Temperature Humidity Index

2.2.1 Comfortability for Human

For the coming February Second ten days, there will be pleasant weather condition over most parts of the country except **Afar, Gambela, Southern-Somali and South Ethiopia** regions, which will be partially (50%) discomfort able for humans day-to-day activity (sign of heat stress expected) as looking in figure 5.




Figure 5: Expected Human comfort

2.2.2 Comfortability for Cattle

Like the human comfortability climate conditions, cattle's comfort index for the next ten days of February Second will have **no significant heat stress** over the country. However, **lowland-border parts** of country will be in mild to moderate heat stress conditions as shown in Figure 6




Figure 6: Expected Dairy (*Top*) and Non-Dairy (*Bottom*) Cattle comfort



3. Summary

Based on the climate-health analysis for February second Dekad, 2025 it has been observed that, there were **no-suitable** climate conditions to the breeding and transmission of malaria in the country. Similarly, over the next 10 days of February 2025 first dekad, there will no-suitable climate condition for the breeding and development mosquitoes except Tips area of north east Tigray.

In terms of climate comfortability conditions, most parts of the country have experienced pleasant conditions for both humans and livestock. Looking ahead to the next 10 days of February 2nd dekad, the low-lying border areas of the country, especially Afar, Gambela, southern Somali, and South Ethiopia region will experience mild (no-significance impact) heat stress, which will affect both humans and cattle.

4. Advisory



Use and implement the following recommendations in places that are favorable for the development of malaria and other vector-borne related diseases;



Attention to any incidence, especially for malaria disease in such favorable areas



Controlling measures and activity are advised



Reducing the environmentally aggravating condition



Awareness creation campaign to the community and sharing of the climate-health update



Avoid any exposure of the community to mosquitoes by ensuring clean environment and using Mosquitoes nets.

