

Ethiopian Meteorology Institute

Health-Meteorology Bulletin

**The product of
Biometeorology and
insurance desk**

Dekadal Bulletin

Volume: 11

No; 31

Issued data: 11/11/2025

Valid until: 30/11/2025

November 2025

**©Ethiopian Meteorology
Institute**

Tell: 251(0)11 6615779

Fax: 251(0)11 6625292

P.O.Box 1090

Website:

<http://www.ethiomet.gov.et/>

Telegram:

<https://t.me/BiometService>



Table of Contents

I. Foreword.....	3
II. Assumptions.....	4
i. Malaria	4
ii. Human heat index:	4
iii. Cattle heat index:	4
1. Weather impact Assessment on Health for First Dekad of November2025	5
1.1 Malaria prone areas during First Dekad of November 2025.....	5
1.2 Climate comfort Conditions.....	5
1.2.1 Comfortability for Human	5
1.2.2 Comfortability for Cattle.....	5
2. Expected Weather Impact on Health for Second dekad of November2025	6
2.1 Expected Mosquito breeding areas	6
2.1 Temperature Humidity Index.....	6
2.2.1 Human Comfort Condition	6
2.2.2 Cattle Comfort Condition	6
4. Advisory.....	7

I. Foreword

This "Climate Information for the Health Sector" Bulletin has been designed to convey essential information regarding the monitoring of human comfort conditions based on the analysis of temperature and humidity data and also for the monitoring of Malaria outbreak areas based on the analysis of temperature and precipitation data. Since the monitoring of temperature and rainfall over a given area can be used to assess the likelihood of outbreak of Malaria with a lag of two months, this information can be an important for early warning tool if used judiciously.

The major objective of this bulletin is in line with the Ethiopia Meteorological Institute strategy of diversifying climate application products to the basic developmental sectors (such as the Health, the water, the agricultural sector etc...). This bulletin can be a very important source of information to Health professionals engaged in the monitoring of Public Health, to Tourism Agents and institutions who advise tourists regarding the comfort conditions of the places to be visited by the tourists and to the researcher who is interested in the field of Bio-Climatology.

We have the opinion that careful and continuous use of this bulletin can benefit to the improvement of early warning and preparedness in the Health sector.

Meanwhile, your comments and constructive suggestions are highly appreciated to make the objective of this bulletin a success,

This same bulletin can be accessed online at: http://www.ethiomet.gov.et/bulletins/health_bulletins

Director General
Ethiopia Meteorology Institute
P.O.Box 1090
Tel: 251(0)11 6615779
FAX 251(0)11 6625292
Web: www.ethiomet.gov.et
Addis Ababa, Ethiopia

II. Assumptions

i. Malaria: According to the International Research Institute for Climate and Society, (IRI), the predicted conditions of rainfall, temperature, and relative humidity are used in determining the degree of incidence for malaria.

- When rainfall is above 80 mm, the temperature is between 25°C and 32°C, and relative humidity is greater than 80%, the region is at high risk and is placed under high incidence.
- When the temperature is between 20°C and 25°C, relative humidity is between 70 and 80%, and rainfall is above 80 mm, then moderate incidence is advised.
- Low incidence for malaria is issued when the temperature is in the range of 18°C-20°C, relative humidity is 60 - 70% and rainfall is above 80 mm.
- No incidence is required when the temperature is less than 18°C, relative humidity less than 60%, and rainfall amount below 80 mm.

Based on these, climate variables have **a one to two months** postponed (delayed) effect on the spread of malaria.

ii. Human heat index: is a measure of how hot it feels when relative humidity is factored with the actual air temperature. The levels of caution for heat index are classified as follows:

- Cold stress when THI is <14, ***Asthma, Pneumonia, Common Cold and flu***
- Comfortable when THI is 14-21, ***pleasanter***
- Moderate when THI is 21-26, ***No more effects***
- Heat stress when THI is >26, ***heat stroke, heat cramps, hyperthermia, respiratory and cardiovascular diseases***

iii. Cattle heat index: The climatic condition for Cattle is a measure that accounts for the combined effects of environmental temperature and relative humidity on cattle. The level of heat stress for cattle classified as follows:

- Not Stressed when THI is <68, ***free from heat stress***
- Stressed threshold when THI is 68 – 71, ***impact less stress starting***
- Mild stress when THI is 72 – 79, ***stress begins and calf rate affected***
- Moderate stress when THI is 80 – 89, ***Milk production affected***
- Severe stress when THI is 90 – 99, ***very significant losses in milk production***
- Extremely stress when THI is >100, ***ultimate dead of cows***



1. Weather impact Assessment on Health for First Dekad of November 2025



1.1 Malaria prone areas during First Dekad of November 2025

Regarding the climatic conditions for malaria breeding and transmission, a low suitable climate condition were observed over Assosa, Illu Aba Bora, all zones of Borena, Guji in Oromia region Southwest Ethiopia region, and Sidama region B during the first dekad of November 2025 as illustrated in Figure 1.




Figure 1: Malaria Prone areas

1.2 Climate comfort Conditions

1.2.1 Comfortability for Human

Over the border parts of Western Amhara, Benishangul Gumuz, Gambela, South Ethiopia region, Southern and Northern Somali and Afar region were experienced heat stress condition for human daily activities during the First dekad of November 2025, But most parts of the country were in pleasant weather conditions as illustrated in Figure 2.




Figure 2: Human comfort index

1.2.2 Comfortability for Cattle

As with human comfort, there were no significant heat stress conditions that affected cattle production; however, as illustrated in Figure 3, some border parts of Afar, Somali, Gambela and Benishangul Gumuz were experienced heat stress conditions of the country.




Figure 3: Cattle Comfort index

2. Expected Weather Impact on Health for Second dekad of November 2025



2.1 Expected Mosquito breeding areas

In the coming Ten days of November 2025, the climate conditions for malaria breeding and transmission will be suitable over, South and North Gondar, Bahir Dar zuria, North Gojam and Awi in Amhara region, Metekel zone in Benishangul Gumuz, Sheka, Bench Maji and Keffa in South West Ethiopia, and East, West and Kellem Wellega, and Illu Aba Bora zones in Oromia regions as illustrated as red in figure 4.




Figure 4: Expected malaria prone areas

2.1 Temperature Humidity Index

2.2.1 Human Comfort Condition

For the coming Second dekad of November 2025, pleasant weather conditions are expected over most parts of the country, except in Afar, South Ethiopia, Southern Somali, **Gambela** and western Amhara (Metema), where conditions will be partially uncomfortable for human daily activities, indicating possible signs of heat stress, as shown in Figure 5.




Figure 5: Expected Human comfort index

2.2.2 Cattle Comfort Condition

Similar to human comfort conditions, cattle are expected to experience mild to moderate heat stress over most lowland areas of Afar, Somali, Gambella, South Ethiopia, Benishangul Gumuz, and western Amhara during the Second dekad of November 2025. However, the western, southern, central, and northern midland and highland areas of the country will be expected to remain free from heat stress, as shown in Figure 6.




Figure 6: Expected Dairy (Top) and Non-Dairy (Bottom) Cattle comfort



3. Summary

According to the first Dekad of November 2025 climate-health analysis, the Western and Southern pocket areas of the country have favourable climate conditions for malaria vector transmission and expansion. As a result, malaria exposure will increase, particularly in Western and Southern Oromia, Assosa, and Southwest Ethiopia. However, heat stress conditions exist in the low-lying border areas of Eastern, Northeaster, Western and Southern Ethiopia, particularly in the Afar, Gambela, Somali, and South Ethiopia regions, affecting human and cattle activity and productivity.

4. Advisory



Use and implement the following recommendations in places that are favourable for the development of malaria and other vector-borne related diseases;

- Attention to any incidence, especially for malaria disease in such favourable areas
- Controlling measures and activity are advised
- Reducing the environmentally aggravating condition
- Awareness creation campaign to the community and sharing of the climate-health update
- Avoid any exposure of the community to mosquitoes by ensuring clean environment and using Mosquitoes nets.

