

Ethiopian Meteorology Institute

Health-Meteorology Bulletin

**The product of
Biometeorology and
insurance desk**

Dekadal Bulletin

Volume: 2

No; 6

Issued data: 01/03/2026

Valid until: 30/04/2026

March 2026

**©Ethiopian Meteorology
Institute**

Tell: 251(0)11 6615779

Fax: 251(0)11 6625292

P.O.Box 1090

Website:

<http://www.ethiomet.gov.et/>

Telegram:

<https://t.me/BiometService>



Table of Contents

I.	Foreword	3
II.	Assumptions	4
	i. Malaria	4
	ii. Human heat index:.....	4
	iii. Cattle heat index:	4
1.	Weather impact Assessment on Health for Second Dekad of February 2026	5
1.1	Malaria prone areas during February 2026	5
1.2	Climate comfort Conditions	5
1.2.1	Comfortability for Human	5
1.2.2	Comfortability for Cattle	5
2.	Expected Weather Impact on Health for first Dekad of March 2026	6
2.1	Expected Mosquito breeding areas	6
2.1	Temperature Humidity Index.....	6
2.2.1	Human Comfort Condition.....	6
2.2.2	Cattle Comfort Condition	6
	Figure 6: Expected.....	6
4.	Advisory	7

I. Foreword

This "Climate Information for the Health Sector" Bulletin has been designed to convey essential information regarding the monitoring of human comfort conditions based on the analysis of temperature and humidity data and also for the monitoring of Malaria outbreak areas based on the analysis of temperature and precipitation data. Since the monitoring of temperature and rainfall over a given area can be used to assess the likelihood of outbreak of Malaria with a lag of two months, this information can be an important for early warning tool if used judiciously.

The major objective of this bulletin is in line with the Ethiopia Meteorological Institute strategy of diversifying climate application products to the basic developmental sectors (such as the Health, the water, the agricultural sector etc...). This bulletin can be a very important source of information to Health professionals engaged in the monitoring of Public Health, to Tourism Agents and institutions who advise tourists regarding the comfort conditions of the places to be visited by the tourists and to the researcher who is interested in the field of Bio-Climatology.

We have the opinion that careful and continuous use of this bulletin can benefit to the improvement of early warning and preparedness in the Health sector.

Meanwhile, your comments and constructive suggestions are highly appreciated to make the objective of this bulletin a success,

This same bulletin can be accessed online at: http://www.ethiomet.gov.et/bulletins/health_bulletins

Director General
Ethiopia Meteorology Institute
P.O.Box 1090
Tel: 251(0)11 6615779
FAX 251(0)11 6625292
Web: www.ethiomet.gov.et
Addis Ababa, Ethiopia

II. Assumptions

i. Malaria: According to the International Research Institute for Climate and Society, (IRI), the predicted conditions of rainfall, temperature, and relative humidity are used in determining the degree of incidence for malaria.

- When rainfall is above 80 mm, the temperature is between 25°C and 32°C, and relative humidity is greater than 80%, the region is at high risk and is placed under high incidence.
- When the temperature is between 20°C and 25°C, relative humidity is between 70 and 80%, and rainfall is above 80 mm, then moderate incidence is advised.
- Low incidence for malaria is issued when the temperature is in the range of 18°C-20°C, relative humidity is 60 - 70% and rainfall is above 80 mm.
- No incidence is required when the temperature is less than 18°C, relative humidity less than 60%, and rainfall amount below 80 mm.

Based on these, climate variables have **a one to two months** postponed (delayed) effect on the spread of malaria.

ii. Human heat index: is a measure of how hot it feels when relative humidity is factored with the actual air temperature. The levels of caution for heat index are classified as follows:

- Cold stress when THI is <14, *Asthma, Pneumonia, Common Cold and flu*
- Comfortable when THI is 14-21, *pleasanter*
- Moderate when THI is 21-26, *No more effects*
- Heat stress when THI is >26, *heat stroke, heat cramps, hyperthermia, respiratory and cardiovascular diseases*

iii. Cattle heat index: The climatic condition for Cattle is a measure that accounts for the combined effects of environmental temperature and relative humidity on cattle. The level of heat stress for cattle classified as follows:

- Not Stressed when THI is <68, *free from heat stress*
- Stressed threshold when THI is 68 – 71, *impact less stress starting*
- Mild stress when THI is 72 – 79, *stress begins and calf rate affected*
- Moderate stress when THI is 80 – 89, *Milk production affected*
- Severe stress when THI is 90 – 99, *very significant losses in milk production*
- Extremely stress when THI is >100, *ultimate dead of cows*



1. Weather impact Assessment on Health for Second Dekad of February 2026



1.1 Malaria prone areas during February 2026

During the February 2026, the prevailing climatic conditions across the country were generally **unfavorable** for malaria vector breeding, development, and transmission dynamics. Temperature and moisture conditions were not conducive for optimal Anopheles mosquito proliferation or for shortening the parasite's extrinsic incubation period, as illustrated in Figure 1.




Figure 1: Malaria Prone areas

1.2 Climate comfort Conditions

1.2.1 Comfortability for Human

During the previous February 2026 month, most parts of the country experienced generally pleasant weather conditions suitable for normal day-to-day human activities. However, moderate to high heat stress conditions were observed over Gambela, South Ethiopia, southern parts of the Somali Region, and Afar Region. These conditions had noticeable impacts on routine human activities, as illustrated in Figure 2.




Figure 2: Human comfort index

1.2.2 Comfortability for Cattle

Similar to the human thermal comfort conditions, **no** significant heat stress that could negatively affect cattle production and productivity was observed during the the February 2026. However, as shown in Figure 3, certain border areas of Afar, Somali, and Gambela regions, experienced noticeable heat stress conditions.




Figure 3: Cattle Comfort index

2. Expected Weather Impact on Health for first Dekad of March 2026



2.1 Expected Mosquito breeding areas

In the upcoming Belg Rainfall benefiting areas, such as South-west Ethiopia, South-Ethiopia, Sidama, Central Ethiopia, Southern and Eastern Oromia, Northshewa (AM), Oromia Special Zone, and western border of Afar regions of the country there will be suitable climate condition for the breeding and transmission of malaria during the coming March 2026, as shown in figure 4.




Figure 4: Expected malaria prone areas

2.1 Temperature Humidity Index

2.2.1 Human Comfort Condition

For the coming First Dekad of March 2026, pleasant weather conditions are expected over most parts of the country, except in Afar, South Ethiopia, Gambela and southern Somali region, where conditions will be partially uncomfortable for human daily activities, indicating possible signs of heat stress, as shown in Figure 5.




Figure 5: Expected Human comfort index

2.2.2 Cattle Comfort Condition

Similar to human comfort conditions, cattle are expected to experience mild to moderate heat stress over most lowland areas of Afar, Somali, Gambella, South Ethiopia, South west Ethiopia, Benishangul Gumuz, Western Amhara and Western Tigray in the coming First Dekad of March 2026. However, the Southern, central, and northern midland and highland areas of the country will be expected to remain free from heat stress, as shown in Figure 6.




Figure 6: Expected Dairy (Top) and Non- Dairy (Bottom) Cattle comfort

3. Summary

4. Advisory



As of February 2026, most parts of the country experienced generally pleasant weather conditions suitable for normal human activities. However, moderate-to-high heat stress was observed and is expected to persist in some lowland areas, particularly Afar, Somali, Gambela, and South Ethiopia, potentially affecting daily activities and vulnerable populations. Climatic conditions during this period were and are expected to remain unfavourable for malaria breeding and transmission countrywide. For livestock, mild to moderate heat stress is expected over lowland areas, including Afar, Somali, Gambela, South Ethiopia, Benishangul-Gumuz, and western Amhara, while midland and highland areas are projected to remain largely free from heat stress.

Use and implement the following recommendations in places that are favourable for the development of malaria and other vector-borne related diseases;

- Strengthen malaria surveillance in favourable areas
- Implement vector control measures
- Promote environmental sanitation
- Conduct community awareness campaigns
- Encourage mosquito net usage
- Monitor heat stress impacts on vulnerable populations and livestock

