

# Ethiopian Meteorology Institute

## Health-Meteorology Bulletin

**The product of  
Biometeorology and  
insurance desk**

**Dekadal Bulletin**

**Volume: 4**

**No; 12**

**Issued data: 01/05/2026**

**Valid until: 30/06/2026**

**May 2026**

**©Ethiopian Meteorology  
Institute**

**Tell: 251(0)11 6615779**

**Fax: 251(0)11 6625292**

**P.O.Box 1090**

**Website:**

<http://www.ethiomet.gov.et/>

**Telegram:**

<https://t.me/BiometService>



## Table of Contents

<b>I.</b>	<b>Foreword</b> .....	3
<b>II.</b>	<b>Assumptions</b> .....	4
1.	Weather impact Assessment on Health for April 2026 .....	5
1.1	Malaria prone areas during April 2026 .....	5
1.2	Climate comfort Conditions .....	5
1.2.1	Comfortability for Human .....	5
1.2.2	Comfortability for Cattle.....	5
2.	Expected Weather Impact on Health for first Dekad of May 2026 .....	6
2.1	Expected Mosquito breeding areas .....	6
2.1	Temperature Humidity Index .....	6
2.2.1	Human Comfort Condition .....	6
2.2.2	Cattle Comfort Condition .....	6
3.	Summary .....	7
4.	Advisory.....	7

## I. Foreword

This "Climate Information for the Health Sector" Bulletin has been designed to convey essential information regarding the monitoring of human comfort conditions based on the analysis of temperature and humidity data and also for the monitoring of Malaria outbreak areas based on the analysis of temperature and precipitation data. Since the monitoring of temperature and rainfall over a given area can be used to assess the likelihood of outbreak of Malaria with a lag of two months, this information can be an important for early warning tool if used judiciously.

The major objective of this bulletin is in line with the Ethiopia Meteorological Institute strategy of diversifying climate application products to the basic developmental sectors (such as the Health, the water, the agricultural sector etc...). This bulletin can be a very important source of information to Health professionals engaged in the monitoring of Public Health, to Tourism Agents and institutions who advise tourists regarding the comfort conditions of the places to be visited by the tourists and to the researcher who is interested in the field of Bio-Climatology.

We have the opinion that careful and continuous use of this bulletin can benefit to the improvement of early warning and preparedness in the Health sector.

Meanwhile, your comments and constructive suggestions are highly appreciated to make the objective of this bulletin a success,

This same bulletin can be accessed online at: [http://www.ethiomet.gov.et/bulletins/health\\_bulletins](http://www.ethiomet.gov.et/bulletins/health_bulletins)

Director General  
Ethiopia Meteorology Institute  
P.O.Box 1090  
Tel: 251(0)11 6615779  
FAX 251(0)11 6625292  
Web: [www.ethiomet.gov.et](http://www.ethiomet.gov.et)  
Addis Ababa, Ethiopia

## II. Assumptions

**i. Malaria:** According to the International Research Institute for Climate and Society, (IRI), the predicted conditions of rainfall, temperature, and relative humidity are used in determining the degree of incidence for malaria.

- When rainfall is above 80 mm, the temperature is between 25°C and 32°C, and relative humidity is greater than 80%, the region is at high risk and is placed under high incidence.
- When the temperature is between 20°C and 25°C, relative humidity is between 70 and 80%, and rainfall is above 80 mm, then moderate incidence is advised.
- Low incidence for malaria is issued when the temperature is in the range of 18°C-20°C, relative humidity is 60 - 70% and rainfall is above 80 mm.
- No incidence is required when the temperature is less than 18°C, relative humidity less than 60%, and rainfall amount below 80 mm.

Based on these, climate variables have **a one to two months** postponed (delayed) effect on the spread of malaria.

**ii. Human heat index:** is a measure of how hot it feels when relative humidity is factored with the actual air temperature. The levels of caution for heat index are classified as follows:

- Cold stress when THI is <14, *Asthma, Pneumonia, Common Cold and flu*
- Comfortable when THI is 14-21, *pleasanter*
- Moderate when THI is 21-26, *No more effects*
- Heat stress when THI is >26, *heat stroke, heat cramps, hyperthermia, respiratory and cardiovascular diseases*

**iii. Cattle heat index:** The climatic condition for Cattle is a measure that accounts for the combined effects of environmental temperature and relative humidity on cattle. The level of heat stress for cattle classified as follows:

- Not Stressed when THI is <68, *free from heat stress*
- Stressed threshold when THI is 68 – 71, *impact less stress starting*
- Mild stress when THI is 72 – 79, *stress begins and calf rate affected*
- Moderate stress when THI is 80 – 89, *Milk production affected*
- Severe stress when THI is 90 – 99, *very significant losses in milk production*
- Extremely stress when THI is >100, *ultimate dead of cows*



# 1. Weather impact Assessment on Health for April 2026

## 1.1 Malaria prone areas during April 2026

During the April 2026, the prevailing climatic conditions across the country were **Low to High** favorable condition for malaria vector breeding and transmission over: all zones western, southern and eastern Oromia regions, all zones of South-west Ethiopia, South Ethiopia, Sidama, Agnuak, Majang, Itang special woredas in Gambela, Metekel and kamashe zones in Benishangul Gumuz, East and west Gajam, Awi, South Gondar, North and South Wollo, WagHamra zones in Amhara, Harari and Dire Dawa city Adm, Daawa, Liben, Afder, Nogob, Erer, Fafan and Siti zones in Somali region as illustrated in Figure 1.




Figure 1: Malaria Prone areas

## 1.2 Climate comfort Conditions

### 1.2.1 Comfortability for Human

During the April 2026, most parts of the country experienced comfortable to moderately a stress weather condition which disrupts human’s day-to-day activities. However, some pocket high land areas of the Northern, Southern, Central and Western parts were in a good condition for human’s day to day activity and all over health status, as illustrated in Figure 2.




Figure 2: Human comfort index

### 1.2.2 Comfortability for Cattle

Similar to the human thermal comfort conditions, **no** significant heat stress that could negatively affect cattle production and productivity were observed during the April 2026, as shown in Figure 3.




Figure 3: Cattle Comfort index



## 2. Expected Weather Impact on Health for first Dekad of May 2026



### 2.1 Expected Mosquito breeding areas

In the upcoming ten day, all zones of Borena, Guji, and Hararge, Bale, and Jimma, Illu Aba Bora, and Kellem Wellega, west Wellega, west Arsi and Arsi in Oromia region, west Gojam and South Wollo zones in Amhara region, all zones of Southwest Ethiopia, South Ethiopia, and Sidama region, Agnuak and Majang zones in Gambela, Liben, Fik, Korahe, Degehabur, Jijiga zones in Somali region will be suitable climate condition for the breeding and transmission of malaria in the coming 1<sup>st</sup> decade of May 2026, as shown in figure 4.




Figure 4: Expected malaria prone

### 2.1 Temperature Humidity Index

#### 2.2.1 Human Comfort Condition

For the coming 1<sup>st</sup> dekad of May 2026, pleasant weather conditions are dominated over most parts of the country. However in Afar, South Ethiopia, Gambela and southern and Northern Somali region, Western Amhara and Benishangul Gumuz region will be partially to fully uncomfortable weather condition prevailing for human daily activities, indicating possible signs of heat stress, as shown in Figure 5.




Figure 5: Expected Human comfort index

#### 2.2.2 Cattle Comfort Condition

None impactful weather condition will be expected to Cattle's product and productivity as well as their health status all over the country in the coming May 1<sup>st</sup> dekad of 2026, as shown in Figure 6.




Figure 6: Expected Dairy (Top) and Non- Dairy (Bottom) Cattle comfort

### 3. Summary

As of April 2026, most parts of the Belg Benefiting areas of the country were experienced favourable climate condition for the breeding and transmission of Malaria. Oromia, Somalie, South Ethiopia, Southwest Ethiopia, Sidama, and central Ethiopia, Gambela, Benishangul Gumuz, Dire Dawa, Harari region were some of the most suitable areas for malaria transmission. However the heat stress condition during this month was not significantly affected both Human and cattle health and productivity.

### 4. Advisory



Use and implement the following recommendations in places that are favourable for the development of malaria and other vector-borne related diseases;

- Strengthen malaria surveillance in favourable areas
- Implement vector control measures
- Promote environmental sanitation
- Conduct community awareness campaigns
- Encourage mosquito net usage
- Monitor heat stress impacts on vulnerable populations and livestock

